

# **Gender Differences in Circadian and Seasonal Variations in Patients with Takotsubo Syndrome- A Multicentre**

## **Registry in Eight University Hospitals in East Japan**

Michiko Yoshizawa<sup>1</sup>, Tomonori Itoh<sup>1,2</sup>, Yoshihiro Morino<sup>1</sup>, Seiichi Tanai<sup>3</sup>, Yuki Ishibashi<sup>4</sup>, Takaaki Komatsu<sup>5</sup>, Isao

Taguchi<sup>5</sup>, Makoto Nishinari<sup>6</sup>, Junya Ako<sup>6</sup>, Hiroyuki Kyono<sup>7</sup>, Taiji Furukawa<sup>7</sup>, Tsutomu Murakami<sup>8</sup>, Yuji Ikari<sup>8</sup>,

Ritsushi Kato<sup>9</sup>, Kazuo Matsumoto<sup>9</sup>, Masashi Sakuma<sup>10</sup>, Hiroyuki Sugimura<sup>11</sup>, Yoshihiro J. Akashi<sup>4</sup>, Hideaki Yoshino<sup>3</sup>

on behalf of the CIRC-8U study group

<sup>1</sup>Division of Cardiology, Department of Internal Medicine, Iwate Medical University, Yahaba, Japan

<sup>2</sup> Division of Community Medicine, Department of Medical Education, Iwate Medical University, Yahaba, Japan

<sup>3</sup>Department of Cardiology, Kyorin University School of Medicine, Tokyo, Japan

<sup>4</sup>Division of Cardiology, Department of Internal Medicine, St. Marianna University School of Medicine, Kawasaki, Japan

<sup>5</sup> Department of Cardiology, Dokkyo Medical University Saitama Medical Center, Koshigaya, Japan

<sup>6</sup>Department of Cardiovascular Medicine, Kitasato University Hospital, Sagamihara, Japan

<sup>7</sup>Department of Medicine, Teikyo University School of Medicine, Tokyo, Japan

<sup>8</sup>Division of Cardiology, Department of Internal Medicine, Tokai University School of Medicine, Isehara, Japan

<sup>9</sup>Division of Cardiology, Saitama Medical University, Moroyama, Japan

<sup>10</sup>Division of Cardiology, Dokkyo Medical University, Mibu, Japan

<sup>11</sup>Division of Cardiology, Dokkyo Medical University Nikko Medical Center, Nikko, Japan

**Corresponding Author:** Tomonori Itoh MD

Tel: 81-19-651-5111

E-mail: tomoitoh@iwate-med.ac.jp

**ORCID**

0000-0001-9277-6567

## **Abstract**

**Objective:** The aim of this study was to clarify the circadian and seasonal variation in addition to identify sex-based differences in Japanese patients with Takotsubo syndrome (TTS). **Methods:** The authors conducted a retrospective observational study to analyse the differences between the groups based on sex. **Patients:** Patients were registered out of each institute registry of the acute coronary syndrome (ACS) which contains a total of 10,622 cases in eight academic hospitals in east Japan. **Results:** Data for 344 consecutive TTS (73 male and 271 female) were extracted from each hospital registry. In-hospital mortality was higher in the male group than in the female group (18% vs 7%;  $p=0.005$ ). With regard to circadian variations in all study patients, TTS events occurred most often in the afternoon and least often during the night. Moreover, the patterns of circadian variations in the female and male groups were the same as that of all study patients. TTS events occurred most frequently in the autumn and least often in the spring in the whole study cohort. Moreover, the seasonal variation in the female group showed the same pattern as that of the whole cohort. However, there was no significant seasonal differences in the incidence of TTS in the male group.

**Conclusions:** In a multicentre study in Japan, seasonal variation was observed in the female group but not in the male group. Circadian variation was observed in both groups. These results suggested that the pathogenesis and clinical features of TTS might be differ based on sex.

**Keywords:** Takotsubo syndrome, circadian variation, seasonal variation, gender differences

## **Introduction**

Takotsubo syndrome (TTS), which is characterized by transient reversible left ventricular dysfunction, was first reported in 1990 in Japan (1). Ninety percent of TTS patients are female and most are over 50 years old (2,3). TTS can occur after several types of stressors (4,5). In general, TTS in female patients is caused by emotional stressors, while TTS in male patients is caused by physical stressors. These patient characteristics have been established in many previous studies (6-8). However, there are few reports regarding circadian and seasonal variations in TTS patients. There is a lack of consensus about circadian (9-13) and seasonal variations in TTS (6,9,12,14), most likely due to the small number of patients. As a result, there are controversies regarding circadian and seasonal variations in patients with TTS. Furthermore, there have been no previous reports about sex-based differences in the circadian or seasonal variations. Given the difference in stressors between male and female TTS patients, there may be differences in chronological variations between those groups of patients. The aims of this study were to elucidate the clinical features of sex-based differences in the circadian and seasonal variations in patients with TTS.

## **Methods**

### *Study Patients*

Patients were registered out of each institute registry of the acute coronary syndrome (ACS) which contains a total of

10,622 cases in the 10 hospitals affiliated with eight medical schools in eastern Japan (the Cardiovascular Research Consortium-8 Universities: CIRC-8U: CIRC-8U: Iwate Medical University, Kitasato University, Tokai University, Dokkyo Medical University, Saitama Medical University, St. Marianna University, Teikyo University, and Kyorin University)). The diagnostic criteria used to diagnose TTS were the Mayo Clinic diagnostic criteria (15). The present study protocol was approved by the ethics committee of each hospital.

#### *Baseline Parameters and Clinical Outcomes*

This was a retrospective observational study that analysed the clinical characteristics of TTS patients. A unified patient registration sheet was used to acquire information such as the clinical history and examination data from the medical records. The types of triggers were classified as suggested by a previous study (5). Emotional triggers included a range of traumatic emotions including grief, interpersonal conflicts, fear, panic, anger, anxiety, and embarrassment. Physical stressors were defined as physical activities, medical conditions, traumatic injury, and surgery.

Cardiac catheterization and echocardiography data were registered in each facility. The left ventricular (LV) ejection fractions (LVEF) of the patients were measured using echocardiography on admission and were evaluated by

the modified Simpson method or Teichholz method. Cardiac catheterization and coronary angiography were performed during the acute phase of admission, mainly on the day of admission. Significant coronary stenosis was defined by visual estimation according to the AHA classification (16). The method of classifying LV asynergy by left ventriculography (LVG) was defined in a previous report (7). LV obstruction was defined as a peak-to-peak pressure gradient greater than 20 mmHg between the LV outflow (LVOT) and the LV apex. Regarding complications during hospitalization, severe pump failure was defined according to the Framingham criteria (17). Cardiogenic shock was defined according to the Killip classification (systolic blood pressure < 90 mmHg or decrease in systolic blood pressure of 30 mmHg, a urine volume  $\leq 20$  cc / hour) and catecholamine use for hypotension.

#### *Definition of Circadian Rhythm and Seasonal Variations*

The time of symptom onset was categorized into four 6-h intervals (night: 12:00 to 5:59 AM; morning: 6:00 to 11:59 AM; afternoon: 12:00 to 5:59 PM; and evening: 6:00 to 11:59 PM) for the circadian analysis. The day of symptom onset was categorized into four 3-month intervals (January to March, April to June, July to September, and October to December).

#### *Statistical Analysis*

Continuous data are presented as the mean $\pm$ standard deviation (SD). Categorical data were analysed with the chi-squared test. The Mann-Whitney U test was used to determine statistically significant differences in non-parametric data between the two groups. Differences were considered significant if the p value was <0.05. The statistical analyses were performed using SPSS for Windows, version 21.0 (Chicago, Illinois, USA). Circadian rhythm and seasonal variations were evaluated by chi-square goodness-of-fit test using StatMate (Version 3.18, ATMS, Tokyo).

## **Results**

### *Comparison between Male and Female Patients*

Out of 10,622 patients with ACS who were admitted to 10 hospitals belonging to CIRC-8U between May 1997 and December 2014, 344 consecutive patients ( $71.6 \pm 11.2$  years of age, 73 male/271 female) were diagnosed with TTS. The clinical characteristics of the study patients are summarized in Table 1. There was no significant difference in age between the sexes. The incidence of smokers was significantly higher in the male group than in the female group, and the incidence of dyslipidaemia was higher in the female group than in the male group. Significant coronary stenosis had a higher prevalence in the male group than in the female group. Significant coronary stenosis was observed in a few patients, but these coronary lesions did not have any haemodynamic effects on the patients.

The incidence of patients with an absence of any stress was 28%, and there was no significant difference in the absence of stress between the groups. However, the incidence of physical stress was significantly higher in the male group than in the female group (64% vs. 46%;  $p=0.007$ ) and the incidence of emotional stress was significantly higher in the female group than in the male group (26% vs. 10%;  $p=0.004$ ). Significant differences between the two groups were not observed in blood pressure or body temperature upon admission to the hospital. However, heart rate, white blood cell count, C-reactive protein and creatinine upon admission were significantly higher in the male group than in the female group. The incidence of significant LV obstruction on cardiac catheterization was 7% in the female group, and no patients in the male group presented. Asynergy of the LVG was categorized as apical type (92%), mid-ventricular type (4%), basal type (1%) and focal type (4%). The proportions of the different LVG classes were not significantly different between the two groups.

#### *In-hospital Complications and Outcomes*

The in-hospital complications and outcomes of the study patients are shown in Table 2. In-hospital mortality was significantly higher in the male group than in the female group (18% vs. 7%;  $p=0.005$ ). The incidence of non-cardiac death was significantly higher in the male group than in the female group (14% vs. 4%;  $p=0.003$ ).



### *Circadian Variations*

Among all study patients, TTS events were most common in the afternoon and least common during the night (Figure 1: A). Moreover, the patterns of circadian variations in the female and male groups were the same as that in the whole study cohort (Figure 1: B: the male group, C: the female group).). Figure of supplemental file showed the circadian rhythm for each trigger. There was less nighttime onset of any trigger. Interestingly, there was also circadian variation in cases with no or unknown stressors. Circadian variation in deaths was as follows: 0.8% at night, 3.2% in the morning, 2.8% during the afternoon, and 2.4% in the evening. Thus, in-hospital mortality was not significantly influenced by onset time ( $p=0.955$ ).

### *Seasonal Variations*

TTS events occurred most often in autumn and least often in spring in whole study cohort (Figure. 2: A). Moreover, the seasonal variation in the female group had the same pattern as that of the whole study cohort. However, there was no significant difference in the incidence of TTS according to season in the male patients (Figure 2: B the male group, C: the female group). In-hospital mortality was not significantly influenced by season as indicated by the following data: 1.5% in spring, 2.9% in summer, 2.3% in autumn, and 2.3% in winter ( $p=0.771$ ).

## Discussion

To the best of the authors' knowledge, this study is the first report of gender differences in both circadian and seasonal variations in patients with TTS. It is noteworthy that all patients in the cohort were of the same ethnicity.

The main results of this study can be summarized as follows. First, this study showed that the in-hospital mortality rate was higher in male patients than in female patients with TTS. Moreover, the cause of death in male patients was more frequently non-cardiac than cardiac, as has been found in previous reports. Second, in the whole study cohort, TTS events were most common in the afternoon and least common during the night. Regarding seasonal variation, the incidence of TTS was highest in the autumn and lowest in the winter and spring in the whole study cohort. Last, the circadian and seasonal variations in the female group were the same as those in the whole study cohort. There was no significant seasonal variation in the male group, although a nearly significant difference according to time of day was observed.

### *Male Patients with Takotsubo Syndrome and Physical Stress*

As has been found in recent studies, in this study, the in-hospital mortality rate was significantly higher in the male group than in the female group (8,18). Moreover, in the male group, the prevalence of physical stress was higher than that of emotional stress (8,19), and the incidence of non-cardiac death was higher than that of cardiac

death (8). The causes of non-cardiovascular death included cancer, infection, sepsis, surgical operation, cerebral haemorrhage, renal failure and gangrene of the foot. White blood cell counts, C-reactive protein and heart rate on admission were higher in the male group than in the female group because the inflammatory reaction is exacerbated by physical stressors. The mortality rate of TTS patients who had another disease before the onset of TTS in the hospital was significantly higher than that of patients without a pre-existing comorbidity (20), and the severity of the comorbidity was thought to be important. Moreover, in-hospital mortality was higher in this study than in the previous study. This is because we assessed data from university hospitals, which possibly treat many numbers of severe cases and patients with underlying diseases than community hospitals. Therefore, the number of physical triggering factors might be higher in our cohort than in those of previous studies.

#### *Circadian Variation of TTS*

Previous reports about circadian variations in patients with TTS have obtained heterogeneous results. There were no circadian variations in patients with TTS in a report from Australia (10). TTS events occurred most often in the morning and least often in the evening in studies from Switzerland (11) and Italy (12), while they were most common in the afternoon and least common in the evening in a study from the United States (13). The reasons for these inconsistent results was that the divisions of time within a day varied, and the numbers of patients in the studies

were small. Moreover, racial differences might affect the results. From Japan in 2007, there was a previous small report on circadian variations that summarized 50 cases of TTS (21). The classification of onset time in that study was similar to that used in our study. There were no significant differences found in circadian variations because of the small number of cases. However, the incidence of onset times during the day was high. The prior study had similar results to those in the present study, which suggests that the onset time of TTS in Japan is most common in the afternoon.

#### *Catecholamine and Circadian Variation*

Catecholamines directly or indirectly affect the brain, vascular system, coronary artery, and myocardium [22]. When people are exposed to stressors, the hypothalamo-pituitary-adrenal axis is activated, and more epinephrine and norepinephrine are released. Estrogens reduce inotropic and chronotropic responses to catecholamines and protect the heart. Postmenopausal depletion may increase catecholamine load [23]. There are still many unclear points about the mechanism of TTS. However, the theory of sympathetic overactivity due to excess catecholamines is considered the most influential. The reason is that pheochromocytoma and acute brain injury are associated with TTS [22], and catecholamine levels in the blood at the acute phase of TTS is higher than that of heart failure caused by ACS [24]. A previous study reported that circadian variations differed between patients with TTS and those with ACS [13]. Serum

catecholamine levels are higher during activity than at rest [25]. Circadian variations in the level of catecholamines might be related to the fact that TTS events are most common in the afternoon. Evaluating the difference in catecholamine levels between the baseline and the peak after exposure to stressors might provide evidence regarding the underlying pathological mechanism and treatment of TTS. Depending on the sub-analysis of our study between circadian variation and TTS trigger, it is also noteworthy that there is circadian variation even in cases without or with unknown stressors. Further study is needed to evaluate between trigger and circadian variation.

#### *Seasonal Variations of TTS*

There have been various reports about seasonal variations in patients with TTS. TTS events occurred most frequently in the summer in the studies from Italy (12) and Germany (22), while a French group reported that the incidence of TTS events was highest in the winter and spring and lowest in the summer (14). Further investigation is needed because the numbers of patients included in the studies from European countries were small. The study of 6837 patients with TTS from the United States reported that TTS occurred most frequently in the summer, with a peak in July, and least frequently in the relatively cold seasons (6). The existing reports are from the Northern Hemisphere, and reports from the Southern Hemisphere are needed. TTS events were reported to be most common in the summer and autumn and least common in the winter and spring in the study of 4306 patients with TTS in Japan, with data

obtained from the diagnosis procedure combination (DPC) (23). The results of study were similar to the results of our study, although there was insufficient diagnostic accuracy in the DPC study. The commonality of results between the two studies suggests that TTS events in Japan occur most frequently in the autumn and least commonly in the winter and spring. The seasonal variation in the female group was similar to that in the whole study cohort. However, there was no significant difference in the incidence of TTS according to season among the male patients. One possible explanation is that TTS events in males are usually caused by physical stress, underlying disease or surgical operations. On the other hand, Japanese women may experience more stressors in autumn, as the stressors associated with human relationships at work and at home peak in the autumn. In another Japanese study, there was significant difference in the proportion of males across seasons (23). The analytic methodology in that study was different from the methodology used in our study. They analysed the changes in the sex ratio of TTS patients according to season. When we applied that analytical method to our data, there were no significant differences in the sex ratio according to season. The seasonal variations of TTS may be affected by not only racial differences but also weather, temperature and atmospheric pressure, so further studies are needed.

### ***Study Limitations***

There are several limitations in this study. First, this study was retrospective. However, the data used in this study was relatively reliable because they were collected from multiple academic hospitals. Second, the number of patients in this study was relatively small. In particular, the number of male patients was small, so the statistical power may be lacking. Another study with a large number of male patients is needed. Third, these data were all obtained from university hospitals, resulting in an institutional bias. It is possible that patients with TTS in general hospitals have different characteristics, so patient data need to be collected from various types of hospitals. Forth, Mayo clinic diagnostic criteria was used in this study. Further study in the future is needed to use new diagnostic criteria. Finally, this study only included data from patients with broken heart syndrome, so it is necessary to collect data from TTS patients with happy heart syndrome (24).

## **Conclusion**

In a multicentre study in Japan, seasonal variations were observed in the female group but not in the male group, while circadian variations were observed in both sexes. These results suggested that the pathogenesis and clinical features of TTS might differ according to sex.

### ***Conflict of interest***

The authors declare that they have no competing interest.

### ***Ethical Approval***

Consent to participate: The study has undergone an ethics review at each participating institution and has been approved.

### ***Informed Consent***

Informed consent was obtained in the form of opt-out on the web-site.

### **Acknowledgments**

The authors are deeply grateful for the support of the following collaborators: Ryohei Sakamoto MD, Satoshi Nakajima MD, Yumiko Okuyama RN and Kayoko Fujiwara (secretary).



## References

1. Dote K, Sato H, Tateishi H, Uchida T, Ishihara M. Myocardial stunning due to simultaneous multivessel coronary spasms: a review of 5 cases. *J Cardiol* 21:203-214, 1991.
2. Patel SM, Chokka RG, Prasad K, Prasad A. Distinctive clinical characteristics according to age and gender in apical ballooning syndrome (takotsubo/stress cardiomyopathy): an analysis focusing on men and young women. *J Card Fail* 19:306-310, 2013.
3. Murakami T, Yoshikawa T, Maekawa Y, et al. Characterization of predictors of in-hospital cardiac complications of takotsubo cardiomyopathy: multi-center registry from Tokyo CCU network. *J Cardiol* 63:269-273, 2014.
4. Ghadri JR, Ruschitzka F, Luscher TF, Templin C. Takotsubo cardiomyopathy: still much more to learn. *Heart* 100:1804-1812, 2014.
5. Ghadri JR, Wittstein IS, Prasad A, et al. International Expert Consensus Document on Takotsubo Syndrome (Part I): Clinical Characteristics, Diagnostic Criteria, and Pathophysiology. *Eur Heart J* 39:2032-2046, 2018.
6. Deshmukh A, Kumar G, Pant S, Rihal C, Murugiah K, Mehta JL. Prevalence of Takotsubo cardiomyopathy in the United States. *Am Heart J* 164:66-71.e61, 2012.
7. Templin C, Ghadri JR, Diekmann J, et al. Clinical Features and Outcomes of Takotsubo (Stress)

- Cardiomyopathy. *N Engl J Med* 373:929-938, 2015.
8. Murakami T, Yoshikawa T, Maekawa Y, et al. Gender differences in patients with takotsubo cardiomyopathy: multi-center registry from Tokyo CCU network. *PLoS One* 10:e0136655, 2015.
  9. Bossone E, Citro R, Eagle KA, Manfredini R. Tako-tsubo cardiomyopathy: is there a preferred time of onset? *Intern Emerg Med* 6:221-226, 2011.
  10. Abdulla I, Kay S, Mussap C, et al. Apical sparing in tako-tsubo cardiomyopathy. *Intern Med J* 36:414-418, 2006.
  11. Eshtehardi P, Koestner SC, Adorjan P, et al. Transient apical ballooning syndrome--clinical characteristics, ballooning pattern, and long-term follow-up in a Swiss population. *Int J Cardiol* 135:370-375, 2009.
  12. Citro R, Previtali M, Bovelli D, et al. Chronobiological patterns of onset of Tako-Tsubo cardiomyopathy: a multicenter Italian study. *J Am Coll Cardiol* 54:180-181, 2009.
  13. Sharkey SW, Lesser JR, Garberich RF, Pink VR, Maron MS, Maron BJ. Comparison of circadian rhythm patterns in Tako-tsubo cardiomyopathy versus ST-segment elevation myocardial infarction. *Am J Cardiol* 110:795-799, 2012.
  14. Mansencal N, El Mahmoud R, Dubourg O. Occurrence of Tako-Tsubo cardiomyopathy and chronobiological variation. *J Am Coll Cardiol* 55:500-501; author reply 501-502, 2010.

15. Prasad A, Lerman A, Rihal CS. Apical ballooning syndrome (Tako-Tsubo or stress cardiomyopathy): a mimic of acute myocardial infarction. *Am Heart J* 155:408-417, 2008.
16. Austen WG, Edwards JE, Frye RL, et al. A reporting system on patients evaluated for coronary artery disease. Report of the Ad Hoc Committee for Grading of Coronary Artery Disease, Council on Cardiovascular Surgery, American Heart Association. *Circulation* 51:5-40, 1975.
17. McKee PA, Castelli WP, McNamara PM, Kannel WB. The natural history of congestive heart failure: the Framingham study. *N Engl J Med* 285:1441-1446, 1971.
18. Brinjkji W, El-Sayed AM, Salka S. In-hospital mortality among patients with takotsubo cardiomyopathy: a study of the National Inpatient Sample 2008 to 2009. *Am Heart J* 164:215-221, 2012.
19. Kurisu S, Inoue I, Kawagoe T, et al. Presentation of tako-tsubo cardiomyopathy in men and women. *Clin Cardiol* 33:42-45, 2010.
20. Isogai T, Yasunaga H, Matsui H, et al. Out-of-hospital versus in-hospital Takotsubo cardiomyopathy: analysis of 3719 patients in the Diagnosis Procedure Combination database in Japan. *Int J Cardiol* 176:413-417, 2014.
21. Kurisu S, Inoue I, Kawagoe T, et al. Circadian variation in the occurrence of tako-tsubo cardiomyopathy: comparison with acute myocardial infarction. *Int J Cardiol* 115:270-271, 2007.
22. Hertting K, Krause K, Harle T, Boczor S, Reimers J, Kuck KH. Transient left ventricular apical ballooning

- in a community hospital in Germany. *Int J Cardiol* 112:282-288, 2006.
23. Isogai T, Matsui H, Tanaka H, Fushimi K, Yasunaga H. Seasonal variation in patient characteristics and in-hospital outcomes of Takotsubo syndrome: a nationwide retrospective cohort study in Japan. *Heart Vessels* 32:1271-1276, 2017.
24. Ghadri JR, Sarcon A, Diekmann J, et al. Happy heart syndrome: role of positive emotional stress in takotsubo syndrome. *Eur Heart J* 37:2823-2829, 2016.

## Figure Captions

**Figure 1: Circadian variation in the study patients:** TTS events were most frequent in the afternoon and least frequent during the night in whole study cohort (A). The pattern of circadian variations in the female and male groups were the same as that in the whole study cohort (B: the male group, C: the female group).

**Figure 2** TTS events occurred most frequently in the autumn and least frequently in the spring in the whole study cohort (A). Seasonal variations in the female group showed the same pattern as that in the whole study cohort. However, there was no significant seasonal variations in the incidence of TTS in the male group (B: the male group, C: the female group)

**Supplemental Figure: The circadian rhythm for each trigger:** There was less nighttime onset of any trigger. Interestingly, there was also circadian variation in cases with no or unknown stressors.

Table 1: Baseline Clinical Characteristics


	All patients (n=344)	Male (n=73)	Female (n=271)	P value
Age (years)	71.6 ± 11.2	71.8 ± 10.4	71.5 ± 11.4	0.899
BMI (kg/m <sup>2</sup> )	21.3 ± 3.9	20.9 ± 3.1	21.5 ± 4.1	0.783
BSA (m <sup>2</sup> )	1.47 ± 0.16	1.61 ± 0.15	1.40 ± 0.23	<0.001
Hypertension	190/332 (57%)	37/71 (52%)	153/261 (59%)	0.346
Dyslipidemia	83/333 (25%)	11/71 (15%)	72/262 (27%)	0.044
Diabetes mellitus	55/332 (17%)	14/70 (20%)	41/262 (16%)	0.371
Current smoking	47/283 (17%)	17/57 (30%)	30/226 (13%)	0.005
Medicines on admission				
β-blockers	19/299 (6%)	6/61 (10%)	13/238 (5%)	0.212
ACE inhibitors/ARB	80/300 (27%)	20/61 (33%)	60/239 (25%)	0.226
Calcium blockers	73/299 (24%)	16/61 (26%)	57/238 (24%)	0.712
Nitrovasodilators	21/303 (7%)	5/62 (8%)	16/241 (7%)	0.693
Symptom				
Chest pain	166/343 (48%)	31/73 (42%)	135/270 (50%)	0.292
No symptom	66/339 (19%)	15/73 (21%)	51/266 (19%)	0.739
Stressors				
Physical stress	162/325 (50%)	45/70 (64%)	117/255 (46%)	0.007
Emotional stress	74/325 (23%)	7/70 (10%)	67/255 (26%)	0.004
Absence of stress	89/325 (27%)	18/70 (26%)	71/255 (28%)	0.764
Vital signs				
Systolic blood pressure (mm Hg)	135.6 ± 31.1	139.2 ± 31.8	134.7 ± 30.9	0.339

Diastolic blood pressure (mm Hg)	79.5 ± 19.0	81.6 ± 21.6	78.9 ± 18.2	0.576
Heart rate (bpm)	94.3 ± 24.9	103.3 ± 28.2	91.9 ± 23.5	0.004
Body temperature (degree)	36.6 ± 1.1	36.9 ± 1.3	36.5 ± 1.1	0.066
Laboratory data				
White blood cell count (cells/ $\mu$ L)	9916 ± 4728	10685 ± 4185	9704 ± 4853	0.011
Peak creatinine kinase (IU/L)	783 ± 2109	799 ± 1838	779 ± 2180	0.065
Peak creatinine kinase MB (IU/L)	30.9 ± 48.3	34.3 ± 56.9	30.0 ± 45.8	0.310
C-reactive protein (mg/dL)	3.3 ± 6.2	5.6 ± 7.1	2.7 ± 5.7	<0.001
Creatinine (mg/dL)	0.9 ± 1.0	1.2 ± 1.1	0.8 ± 0.9	<0.001
ST elevation	208/286 (73%)	42/57 (74%)	166/229 (72%)	1.000
LVEF (%) by TTE	45.9 ± 13.0	44.7 ± 13.2	46.2 ± 13.0	0.544
LVEF (%) by left ventriculography	46.6 ± 11.8	44.5 ± 11.7	47.0 ± 11.8	0.122
Coronary stenosis by cardiac catheterization or coronary angiography	23/285 (8%)	11/58 (19%)	12/227 (5%)	0.002
Cardiac catheterization				
Apical type	227/248 (91.5%)	44/47 (93.6%)	183/201 (91.0%)	0.539
Midventricular type	9/248 (3.6%)	1/47 (2.1%)	8/201(4.0%)	
Basal type	2/248 (0.8%)	1/47 (2.1%)	1/201 (0.5%)	
Focal type	10/248 (4.0%)	1/47 (2.1%)	9/201 (4.5%)	
LV obstruction	10/185 (5%)	0/30 (0%)	10/155 (6%)	0.162


Table 2: In-hospital Complications and Outcome

	All patients (n=344)	Male (n=73)	Female (n=271)	P value
Severe heart failure	102/341 (30%)	25/73 (34%)	77/268 (29%)	0.388
Cardiogenic shock	50/340 (15%)	14/73 (19%)	36/267 (13%)	0.262
Arrhythmia				
Atrial fibrillation	33/340 (10%)	7/73 (10%)	26/267 (10%)	1.000
Atrial tachycardia				
Ventricular tachycardia	14/340 (4%)	4/73 (5%)	10/267 (4%)	0.510
Ventricular fibrillation				
Complete atrioventricular block	7/340 (2%)	1/73 (1%)	6/267 (2%)	1.000
Sick Sinus Syndrome				
Left ventricular thrombus	6/336 (2%)	0/72 (0%)	6/264 (2%)	0.581
Recurrence	4/355 (1%)	1/72 (1%)	3/263 (1%)	1.000
Death	31/341 (9%)	13/71 (18%)	18/270 (7%)	0.005
Cardiovascular death	11/341 (3%)	3/71 (4%)	8/270 (3%)	0.704
Others	20/341 (6%)	10/71 (14%)	10/270 (4%)	0.003






**Total (n=251)**




**Male group (n=54)**




**Female group (n=197)**


Figure 1



**Total (n=342)**



**Male group (n=72)**



**Female group (n=270)**

Figure 2